

# Dosing for Infants and Children

From Your Healthcare Professional



**DOSE: Every 4 hours as needed. DO NOT GIVE MORE THAN 5 DOSES IN 24 HOURS.**

If possible, use weight to dose; otherwise, use age.

mL = milliliter



## Infants' TYLENOL® Oral Suspension

**Active Ingredient:**  
Acetaminophen 160 mg  
(in each 5 mL)

Available in:



Use product only as directed.



## Children's TYLENOL® Oral Suspension

**Active Ingredient:**  
Acetaminophen 160 mg  
(in each 5 mL)

Available in:



Use product only as directed.

WEIGHT	AGE	Infants' TYLENOL®	Children's TYLENOL®
6-11 POUNDS	0-3 MONTHS	1.25 mL	—
12-17 POUNDS	4-11 MONTHS	2.5 mL	—
18-23 POUNDS	12-23 MONTHS	3.75 mL	—
24-35 POUNDS	2-3 YEARS	5 mL	5 mL
36-47 POUNDS	4-5 YEARS	—	7.5 mL
48-59 POUNDS	6-8 YEARS	—	10 mL
60-71 POUNDS	9-10 YEARS	—	12.5 mL
72-95 POUNDS	11 YEARS	—	15 mL

## Kids' Wellness Tracker



This free, all-in-one tool from the makers of Children's TYLENOL® and Children's MOTRIN® makes it easy to track your child's height, weight, BMI, vaccines, symptoms, and medicines, plus calculate dosing.

Available on the App Store™ and the Google Play Store™



Google Play is a trademark of Google Inc. App Store is a service mark of Apple Inc. Standard download rates may apply.

## IMPORTANT INSTRUCTIONS for Proper Use

Today's Date: \_\_\_\_\_

This dosing recommendation from your doctor will expire in 14 DAYS.

- Read and follow the label on all TYLENOL® products.
- Repeat dose every 4 hours while symptoms last.
- Do NOT give more than 5 doses in 24 hours.
- Do NOT use with any other product containing acetaminophen.
- Use only the dosing device (syringe or dosing cup) that came with the product.
- **All Infants' TYLENOL® and Children's TYLENOL® Oral Suspension products have the same acetaminophen concentration (160 mg/5 mL).**

# Dosing for Infants and Children

From Your Healthcare Professional

Concentrated  
**MOTRIN**  
Infants' Drops

**Children's  
MOTRIN**

**DOSE: Every 6-8 hours as needed. DO NOT GIVE MORE THAN 4 DOSES IN 24 HOURS.**

If possible, use weight to dose; otherwise, use age.

mL = milliliter



## Infants' MOTRIN® Concentrated Drops

**Active Ingredient:**  
Ibuprofen 50 mg (NSAID)\*  
(in each 1.25 mL)

\*Nonsteroidal anti-inflammatory drug  
Use product only as directed.

**Available in:**  
Berry  
Dye-Free Berry










## Children's MOTRIN® Oral Suspension

**Active Ingredient:**  
Ibuprofen 100 mg (NSAID)\*  
(in each 5 mL)

\*Nonsteroidal anti-inflammatory drug  
Use product only as directed.

**Available in:**  
Bubble Gum  
Grape  
Berry  
Dye-Free Berry

WEIGHT	AGE	Infants' MOTRIN® Concentrated Drops	Children's MOTRIN® Oral Suspension
—	<6 MONTHS	—	—
12-17 POUNDS	6-11 MONTHS	1.25 mL 	—
18-23 POUNDS	12-23 MONTHS	1.875 mL 	—
24-35 POUNDS	2-3 YEARS	—	5 mL 
36-47 POUNDS	4-5 YEARS	—	7.5 mL 
48-59 POUNDS	6-8 YEARS	—	10 mL 
60-71 POUNDS	9-10 YEARS	—	12.5 mL 
72-95 POUNDS	11 YEARS	—	15 mL 

## Kids' Wellness Tracker



This free, all-in-one tool from the makers of Children's TYLENOL® and Children's MOTRIN® makes it easy to track your child's height, weight, BMI, vaccines, symptoms, and medicines, plus calculate dosing.

Available on the App Store<sup>SM</sup> and the Google Play Store<sup>TM</sup>



Google Play is a trademark of Google Inc. App Store is a service mark of Apple Inc. Standard download rates may apply.

## IMPORTANT INSTRUCTIONS for Proper Use

Today's Date: \_\_\_\_\_

This dosing recommendation from your doctor will expire in 14 DAYS.

- Read and follow the label on all MOTRIN® products.
- Repeat dose every 6-8 hours as needed.
- Do NOT give more than 4 doses in 24 hours.
- Use only the dosing device (syringe or dosing cup) that came with the product.